

CASE REPORT

Open Access



To reperfuse or not to reperfuse: a case report of Wellens' syndrome with suspected COVID-19 infection

I Gde Rurus Suryawan^{1*†}, Jordan Bakhriansyah^{2,3†}, Mia Puspitasari³, Parama Gandhi¹, Ryan Enast Intan¹ and Firas Farisi Alkaff⁴

Abstract

Background: Wellens' syndrome is known to be associated with left anterior descending artery occlusion that could lead to an extensive anterior wall myocardial infarction. Thus, emergency cardiac catheterization is needed. However, during coronavirus disease 2019 (COVID-19) pandemic, it is recommended for hemodynamically stable acute coronary syndrome patients with COVID-19 infection to be treated conservatively in an isolated hospital ward.

Case presentation: We report an 85-year-old patient with chief complaints of typical, squeezing chest pain in the past 4 h. The patient had a high fever, dyspnea, sore throat, and fatigue for 3 days. He had previously come into contact with COVID-19 positive relatives. The patient was hemodynamically stable and pulmonary auscultation revealed coarse rales in the entire lung. Electrocardiography (ECG) evaluation during the pain episode showed non-specific ST-T changes in lead V2-V5. After sublingual nitrate was administered, ECG evaluation during the pain-free period revealed a biphasic T wave inversion in lead V2 and V3. Laboratory workup showed elevated cardiac marker and leucopenia with neutrophilia and lymphopenia. Rapid immunochromatographic test and initial severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) reverse transcription-polymerase chain reaction (RT-PCR) evaluation from nasopharyngeal swab showed negative results. However, radiographic evaluations suggest the diagnosis of COVID-19 infection. While waiting for the second RT-PCR evaluation, the patient was diagnosed with Wellens' syndrome with suspected COVID-19 infection. The patient was treated conservatively according to national guidelines and scheduled for elective cardiac catheterization. On the third day, the patient felt better and insisted on being discharged home. Ten days after discharged, the patient died of myocardial infarction.

Conclusion: Emergency cardiac catheterization should be done for patient with Wellens' syndrome, regardless of the COVID-19 infection status.

Keywords: Cardiac catheterization, Case reports, COVID-19, Myocardial infarctions

* Correspondence: igr.suryawan@gmail.com

†I. Gde Rurus Suryawan and Jordan Bakhriansyah contributed equally to this work.

¹Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Jl. Mayjen Prof. Dr. Moestopo No. 6-8, Surabaya, East Java 60286, Indonesia
Full list of author information is available at the end of the article



© The Author(s). 2020 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

Background

Wellens' syndrome was first reported in 1982 and is known to be associated with left anterior descending (LAD) artery occlusion. If left untreated, it could lead to an extensive anterior wall myocardial infarction [1]. A later study with 18 months follow-up period showed that the mortality rate was remarkably high in patients treated conservatively compared to patients treated with cardiac catheterization (26.67% vs. 0.88%) [2]. Thus, emergency cardiac catheterization is warranted in patients presenting with this syndrome.

However, during this coronavirus disease 2019 (COVID-19) pandemic, more caution is taken into consideration for all invasive procedures, including for the cardiac catheterization procedure. In patients with COVID-19 infection, the balance of staff exposure and patient benefit should be weighed carefully. Based on the patient's risk, conservative therapy may be sufficient for non-ST-segment elevation myocardial infarction (NSTEMI) patient with COVID-19 [3]. The Indonesian Heart Association also published a national practical clinical guideline for NSTEMI patient with COVID-19. It is recommended that patients with stable hemodynamic to be treated conservatively in isolated hospital ward [4]. In this report, we present a patient with Wellens' syndrome with suspected COVID-19 infection based on the clinical symptoms and radiographic findings. The patient was treated conservatively according to the national guideline for NSTEMI with COVID-19 infection.

Case presentation

An 85-year-old man came to the emergency room with the chief complaint of typical, squeezing chest pain in

the past 4 h. The patient also experienced diaphoresis and nausea following chest pain. In the past 3 days, the patient had a high fever, dyspnea, sore throat, and fatigue. Past medical history of type 2 diabetes mellitus or hypertension was denied. He had a history of contact with one of his relatives who tested positive for severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) based on reverse transcription-polymerase chain reaction (RT-PCR) evaluation.

Vital signs on admission were as follows: blood pressure 130/90 mmHg, respiratory rates 26 times/min, heart rate 104 beats/min, right axillary temperature 39 °C, oxygen saturation 94% at room air, and became 99% with the simple mask with 6 L/min oxygen. Pulmonary auscultation revealed coarse rales in the entire lung. Other physical examinations were within normal limit. Twelve-lead electrocardiography (ECG) performed when the patient was in pain showed non-specific ST-T changes in lead V2-V5 (Fig. 1a). After receiving sublingual nitrate, the chest pain subsided, and the ECG evaluation showed biphasic T wave inversion and minimally elevated ST-segment in lead V2 and V3 (Fig. 1b). Before the patient was transferred to the hospital ward, the ECG evaluation in pain-free period revealed deeply inverted T waves in lead V2-V4 (Fig. 1c).

Laboratory evaluation revealed leucopenia ($3.88 \times 10^3/\mu\text{l}$) with neutrophilia (89.4%) and lymphopenia (3.6%), thrombocytopenia ($102 \times 10^3/\mu\text{l}$), elevated aspartate transaminase (AST) (80.3 U/L), and slightly elevated alanine aminotransferase (ALT) (44.4 U/L). Creatine kinase myocardial band (CK-MB) was also increased (10.4 ng/mL). Serum creatinine and blood glass analysis were within normal limits. The Global Registry of Acute




Fig. 1 Electrocardiography evaluation (a) performed on admission in the emergency room during pain period showed non-specific ST-T changes in lead V2-V5, (b) performed on the painless period after sublingual nitrate showed biphasic T waves inversions and minimally elevated ST segment in lead V2 and V3, (c) performed before transferred to hospital ward during the painless period showed deeply inverted T waves in leads V2-V4




Fig. 2 Chest X-ray showed preceding consolidation persisted with new consolidative changes in the left apical-middle-lower zone and the right lower peripheral region

Coronary Events (GRACE) score was 159 and CRUS ADE bleeding score was 37. Chest X-ray showed preceding consolidation persisted with new consolidative changes in the left apical-middle-lower zone and the right lower peripheral region (Fig. 2). Chest computed tomography scan (CT scan) revealed diffuse pneumonia in both lungs with multifocal ground-glass opacities and crazy paving patterns (Fig. 3), a common finding in patients with COVID-19 infection. However, SARS-CoV-2 rapid immunochromatographic test (Wondfo Biotech, Guangzhou, China) showed a non-reactive result. The




Fig. 3 Chest computed tomography scan showed diffuse pneumonia in both lungs with multifocal ground-glass opacities and crazy paving patterns

initial RT-PCR assay (Abbott RealTime SARS-CoV-2 assay, Abbott Molecular Inc., Illinois, USA) from nasopharyngeal swab was then performed, but it also showed a negative result.

Since the diagnosis of COVID-19 infection could not be ruled out until the RT-PCR assay was repeated, the patient was diagnosed with Wellens' syndrome with suspected COVID-19 infection. Because the patient was categorized as high-risk NSTEMI (high GRACE score but with stable hemodynamic) with high neutrophil-to-lymphocyte ratio and suspected with COVID-19 infection, the patient was treated conservatively in the intensive care unit (ICU) isolation ward while waiting for the early elective cardiac catheterization. The patient received double antiplatelet therapy (DAPT) of aspirin (80 mg once daily) and clopidogrel (75 mg once daily), fondaparinux (2.5 mg once daily), atorvastatin (80 mg once daily), bisoprolol (2.5 mg once daily), isosorbide dinitrate pump (1 mg per hour), paracetamol (500 mg thrice daily), and methisoprinol (500 mg thrice daily).

On the third day, the patient's oxygen saturation was 98% without oxygen supplementation, and ECG evaluation reverts to biphasic T wave in lead V2 and V3 (Fig. 4). CK-MB level was still above the normal limit (19 ng/mL). The patient insisted on being discharged and refused to be referred for early elective cardiac catheterization because he already felt better. The patient and his family signed the consent form to be discharged home despite the high chance of myocardial infarction in the near future. The patient was also aware that the diagnosis of COVID-19 infection could not be ruled out yet because second RT-PCR from nasopharyngeal swab had not been performed yet, thus he and his family had to self-quarantine at home for 14 days.

On the fourth day, the patient was discharged and received aspirin (80 mg once daily) and clopidogrel (75 mg once daily) for his take-home medicine. Two weeks later, in a follow-up session via telephone, one of the family members informed that the patient already died 10 days after being discharged from our hospital due to cardiac arrest secondary to new-onset ST-elevation myocardial infarction. Due to the limited facilities in the other hospital, the patient did not undergo coronary angiography.

Discussion

Our patient has fulfilled the suspect case criteria for COVID-19 by WHO guideline as followed: A patient with acute respiratory illness (fever and at least one sign/symptom of respiratory disease, e.g., cough, shortness of breath), AND a history of travel to or residence in a location reporting community transmission of COVID-19 disease or having been in contact with a confirmed or probable COVID-19 case during the 14 days prior to symptom onset or patients with severe acute respiratory illness (fever and at least one sign/symptom of




Fig. 4 Electrocardiography re-evaluation on the third day showed biphasic T waves in lead V2 and V3

respiratory disease, e.g., cough, shortness of breath; AND requiring hospitalization) AND in the absence of an alternative diagnosis that thoroughly explains the clinical presentation [5]. This was also supported by the laboratory abnormalities found in our patient, which were lymphopenia, leucopenia, thrombocytopenia, elevated creatinine, AST and ALT, and hypoxemia from blood gas analysis, and also pneumonia by chest X-ray and CT scan findings (bilateral, peripheral, patchy opacities on chest X-ray and bilateral ground-glass opacities, crazy paving, and multifocal consolidation from chest CT scan) that suggest high probability for COVID-19 infection. It is suggested that findings from chest CT scans usually peak around 9–13 days [6, 7].

WHO criteria for confirmed COVID-19 was based on the detection of unique sequences of virus SARS-CoV-2 RNA by nucleic acid amplification tests such as real-time RT-PCR and needed at least two positive results [8]. For initial diagnostic testing, Centers for Disease Control and Prevention (CDC) recommends collecting and testing an upper respiratory specimen with a nasopharyngeal swab as the preferred specimen choice [9]. However, multiple negative tests are required to exclude a diagnosis of COVID-19.

CDC also stated that negative SARS-CoV-2 results from RT-PCR do not preclude COVID-19 infection and should not be used as the sole basis for patient treatment decisions, especially when it is not supported with the clinical observations, patient history, and epidemiological information [9]. It is because RT-PCR has the sensitivity as low as 6–70% for initial diagnosis despite its high specificity [10]. In our case, although the initial SARS-CoV-2 RT-PCR showed a negative result, the chest CT scan showed a typical manifestation of COVID-19. This might be explained by the findings from previous study that the sensitivity of the initial

chest CT scan is greater than the initial RT-PCR assay (98% vs 71%, $p < 0.001$) [11].

To establish the diagnosis of Wellens' syndrome, it is suggested that several criteria be fulfilled, which includes (1) deep symmetrically inverted T waves or biphasic T waves in lead V2 and V3, (2) isoelectric or minimally elevated (< 1 mm) ST-segment, (3) absence of precordial Q waves, (4) history of angina, (5) pattern present during pain-free period, and (6) normal or mildly elevated creatine phosphokinase (less than two times normal upper limit) [12]. In our case, the patient fulfilled all criteria for Wellens' syndrome except the cardiac marker. However, since the cardiac marker is known to be frequently abnormal in patients with COVID-19 [13], we argued that the cardiac marker criterion could be exempted in this situation.

There were some reports regarding the association between COVID-19 infection and cardiovascular complications including myocardial injury, myocarditis, deep vein thrombosis (DVT), and pulmonary embolism (PE) [14]. Our case might be correlated to COVID-19 infection-induced myocardial injury, infarction, or inflammation due to systemic inflammation response, marked by an elevated CK-MB level. However, it is unlikely that our patient had DVT because there were no supporting clinical findings such as warmth or pain in the extremity or asymmetrical swelling [15]. PE could also be ruled out because there was also no filling defect in the pulmonary artery in the chest CT scan evaluation [16].

It could be argued that this type of case is usually diagnosed as high-risk anterior NSTEMI. However, we would like to stress out the use of Wellens' syndrome nomenclature to underline the high probability of total or near-total LAD occlusion that is not commonly found in high-risk NSTEMI patients. Patients with Wellens' syndrome will develop extensive anterior wall infarction

if aggressive intervention is not undertaken, despite the relief of symptoms with medical management. Half of the patients will develop the infarction within 1 week after the admission [1]. Thus, in normal situation, our patient should have undergone emergency cardiac catheterization. Other than that, our patient also had a GRACE score of 159. European Society of Cardiology (ESC) and American College of Cardiology/American Heart Association recommend an invasive strategy should be performed in less than 24 h in patient with high-risk NSTEMI (GRACE score more than 140) [17, 18].

However, in patients with suspected COVID-19 infection, the algorithm management is different. National guideline published by the Indonesian Heart Association recommends conservative treatment in the isolated hospital ward if the patients have stable hemodynamic to reduce transmission risk of COVID-19, especially when a special standardized facility is not available [4]. This recommendation was in line with the Chinese Society of Cardiology guideline that recommends patients with high-risk NSTEMI to be hospitalized and treated conservatively in designated hospital [19]. American College of Cardiology suggests that in patient with stable NSTEMI, conservative therapy may be sufficient on the basis of patient risk [3]. In contrary, guideline published by ESC recommends patients with high-risk NSTEMI to still be treated with an early invasive strategy in less than 24 h after admission in COVID-19 designated hospital [20]. According to the Egyptian Society of Cardiology guidelines, patients with high-risk NSTEMI should undergo early catheterization in less than 24 h. However, it is only possible if the hospital is not overwhelmed and all the precautions to prevent the dissemination of infection and protect the medical staff are adopted. Nevertheless, if the prevalence of COVID-19 increases and causes overburden of the health system resources, patients with high-risk NSTEMI should be hospitalized and treated conservatively in isolation wards or ICU in non-designated hospital [21].

According to the national guideline, DAPT (clopidogrel or ticagrelor and aspirin) and high-dose statin should be given for conservative treatment during hospitalization [4]. We opted to treat our patient with clopidogrel because of the moderate bleeding risk (CRUSADE score 37). Recent meta-analysis study showed that ticagrelor was associated with a higher risk of major bleeding compared to clopidogrel in East Asian patients with acute coronary syndrome [22]. In addition to those recommended treatment, the patient also received fondaparinux. After the hospitalization, the patient had been given DAPT for take-home medicine. It is recommended that DAPT should be given for 1 year after discharged home [17].

The limitations of this report were the absence of coronary angiography and echocardiography evaluation. Therefore, the diagnosis of the patients could not be confirmed and the differential diagnosis such as PE and myocarditis could not be totally excluded. The coronary angiography was not performed because it was not recommended by the Indonesian Heart Association. Echocardiography evaluation was not performed because there was no published guideline in performing echocardiography to patients with suspected COVID-19 infection and there was also a shortage of standardized personal protective equipment when this case report occurred.

Conclusion

In the case of acute coronary syndrome in the COVID-19 pandemic situation where the risk of infectious spread is very high, risk stratification is essential to determine the treatment strategy. Following the national guideline in this situation, high-risk NSTEMI with conservative management is preferred for treatment in the acute phase, with favorable outcomes in the acute phase. However, in the case of Wellens' syndrome, where significant LAD occlusion is suspected, urgent early cardiac catheterization should be done, regardless of the COVID-19 infection status. Recognition of the ECG pattern of Wellens' syndrome is also crucial because Wellens' syndrome has a poor prognosis despite the lack of symptoms in early stable conditions.

Abbreviations

ALT: Alanine Aminotransferase; AST: Aspartate Transaminase; CDC: Centers for Disease Control and Prevention; CK-MB: Creatine kinase myocardial band; COVID-19: Coronavirus disease 2019; CT scan: Computed tomography scan; DAPT: Double anti platelet therapy; ECG: Electrocardiography; ESC: European Society of Cardiology; GRACE: Global Registry of Acute Coronary Events; ICU: Intensive care unit; LAD: Left anterior descending; NSTEMI: Non-ST-segment elevation myocardial infarction; RT-PCR: Reverse transcription polymerase chain reaction; SARS-CoV-2: Severe acute respiratory syndrome coronavirus 2

Acknowledgements

The authors would like to thank the Institute of Tropical Disease, Universitas Airlangga, Surabaya, Indonesia, for performing the rapid immunochromatographic test and RT-PCR assay of the patient.

Authors' contributions

IGRS and JB contribute equally to the study design, data analysis and interpretation, and substantially revised the manuscript. MP contributes to the data analysis and interpretation. PG contributes to the data collection. REI and FFA contribute in literature review of the case and drafting the manuscript. All authors have read and approved the final manuscript.

Funding

No funding was received.

Availability of data and materials

Available upon request.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Written informed consent was obtained from the patient for publication of this case report and accompanying images.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Airlangga – Dr. Soetomo General Academic Hospital, Jl. Mayjen Prof. Dr. Moestopo No. 6-8, Surabaya, East Java 60286, Indonesia.

²Department of Cardiology and Vascular Medicine, Husada Utama Hospital, Surabaya, Indonesia. ³Faculty of Medicine, Universitas Surabaya, Surabaya, Indonesia. ⁴Department of Pharmacology, Faculty of Medicine Universitas Airlangga, Surabaya, Indonesia.

Received: 18 May 2020 Accepted: 27 August 2020

Published online: 09 September 2020

References

- de Zwaan C, Bar FW, Wellens HJ (1982) Characteristic electrocardiographic pattern indicating a critical stenosis high in left anterior descending coronary artery in patients admitted because of impending myocardial infarction. *Am Heart J* 103(4 Pt 2):730–736
- de Zwaan C, Bar FW, Janssen JH, Cheriex EC, Dassen WR, Brugada P et al (1989) Angiographic and clinical characteristics of patients with unstable angina showing an ECG pattern indicating critical narrowing of the proximal LAD coronary artery. *Am Heart J* 117(3):657–665
- Welt FGP, Shah PB, Aronow HD, Bortnick AE, Henry TD, Sherwood MW et al (2020) Catheterization laboratory considerations during the coronavirus (COVID-19) pandemic: from ACC's Interventional Council and SCAI. *J Am Coll Cardiol*
- Indonesian Heart Association (2020) Clinical Practical Guideline Indonesian Heart Association. NSTEMI with COVID-19 suspicion. Pengurus Pusat Perhimpunan Dokter Spesialis Kardiovaskular Indonesia, Jakarta
- World Health Organization. Global surveillance for COVID-19 caused by human infection with COVID-19 virus. 2020.
- Xie X, Zhong Z, Zhao W, Zheng C, Wang F, Liu J (2020) Chest CT for typical 2019-nCoV pneumonia: relationship to negative RT-PCR testing. *Radiology*. 200343
- Jamil S, Mark N, Carlos G, Dela Cruz CS, Gross JE, Pasnick S (2020) Diagnosis and management of COVID-19 disease. *Am J Respir Crit Care Med*
- World Health Organization. Laboratory testing for 2019 novel coronavirus (2019-nCoV) in suspected human cases. 2020.
- Centers for Disease Control and Prevention. CDC diagnostic test for COVID-19 2020 [updated 14 April 2020; cited 18 April 2020]. Available from: https://www.cdc.gov/coronavirus/2019-ncov/php/testing.html?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Fcoronavirus%2F2019-ncov%2Fabout%2Ftesting.html.
- Ai T, Yang Z, Hou H, Zhan C, Chen C, Lv W et al (2020) Correlation of chest CT and RT-PCR testing in coronavirus disease 2019 (COVID-19) in China: a report of 1014 cases. *Radiology*. 200642
- Fang Y, Zhang H, Xie J, Lin M, Ying L, Pang P et al (2020) Sensitivity of chest CT for COVID-19: comparison to RT-PCR. *Radiology*. 200432
- Rhinehardt J, Brady WJ, Perron AD, Mattu A (2002) Electrocardiographic manifestations of Wellens' syndrome. *Am J Emerg Med* 20(7):638–643
- Bansal M (2020) Cardiovascular disease and COVID-19. *Diabet Metabolic Syndrome* 14(3):247–250
- Long B, Brady WJ, Koyfman A, Gottlieb M (2020) Cardiovascular complications in COVID-19. *Am J Emerg Med* 38(7):1504–1507
- Stone J, Hangge P, Albadawi H, Wallace A, Shamoun F, Knutti MG et al (2017) Deep vein thrombosis: pathogenesis, diagnosis, and medical management. *Cardiovasc Diagnosis Ther* 7(Suppl 3):S276–S284
- Liu M, Tao XC, Zhai Z, Ma Z, Zhu L, Luo J (2020) The filling defect of pulmonary artery, an imaging finding what we should know. *Pulmonary Circ* 10(1):2045894020910687
- Roffi M, Patrono C, Collet JP, Mueller C, Valgimigli M, Andreotti F et al (2016) 2015 ESC guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation: task force for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation of the European Society of Cardiology (ESC). *Eur Heart J* 37(3):267–315
- Amsterdam EA, Wenger NK, Brindis RG, Casey DE Jr, Ganiats TG, Holmes DR Jr et al (2014) 2014 AHA/ACC guideline for the management of patients with non-ST-elevation acute coronary syndromes: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *J Am Coll Cardiol* 64(24):e139–e228
- Han Y, Zeng H, Jiang H, Yang Y, Yuan Z, Cheng X et al (2020) CSC expert consensus on principles of clinical management of patients with severe emergent cardiovascular diseases during the COVID-19 epidemic. *Circulation*.
- The European Society for Cardiology. ESC guidance for the diagnosis and management of CV disease during the COVID-19 pandemic 2020 [updated 10 June 2020]. Available from: <https://www.escardio.org/Education/COVID-19-and-Cardiology/ESCCOVID-19-Guidance>.
- Shaheen S, Awwad O, Shokry K, Abdel-Hamid M, El-Etriby A, Hasan-Ali H et al (2020) Rapid guide to the management of cardiac patients during the COVID-19 pandemic in Egypt: "a position statement of the Egyptian Society of Cardiology". *Egyptian Heart J* 72(1):30
- Misumida N, Aoi S, Kim SM, Ziada KM, Abdel-Latif A (2018) Ticagrelor versus clopidogrel in East Asian patients with acute coronary syndrome: systematic review and meta-analysis. *Cardiovasc Revasc Med* 19(6):689–694

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](https://www.springeropen.com)